

Table 3

(From Journal of Nuclear Medicine, Volume 41, pages 863-873, 2000).

Summary of Recommendations for Radiopharmaceuticals Excreted in the Breast Milk

<u>Pharmaceutical</u>	<u>Administered Activity, mCi (MBq)</u>	<u>Counseling?*</u>	<u>Advisory</u>
Ga-67 Citrate	5.0 (185)	Yes	Cessation
Tc-99m DTPA	20 (740)	No	
Tc-99m MAA	4 (148)	Yes	12 hr
Tc-99m Pertechnetate	5 (185)	Yes	4 hr
I-131 NaI	150 (5550)	Yes	Cessation
Cr-51 EDTA	0.05 (1.85)	No	
Tc-99m DISIDA	8 (300)	No	
Tc-99m glucoheptonate	20 (740)	No	
Tc-99m HAM	8 (300)	No	
Tc-99m MIBI	30 (1110)	No	
Tc-99m MDP	20 (740)	No	
Tc-99m PYP	20 (740)	No	
Tc-99m RBC's in vivo	20 (740)	Yes	12 hr
Tc-99m RBC's in vitro	20(740)	No	
Tc-99m Sulfur Colloid	12 (444)	No	
In-111 WBC's	0.5 (18.5)	No	
I-123 NaI	0.4 (14.8)	Yes	Cessation
I-123 OIH	2 (74)	No	
I-123 MIBG	10 (370)	Yes	48 hr
I-125 OIH	0.01 (0.37)	No	
I-131 OIH	0.3 (11.1)	No	
Tc-99m DTPA Aerosol	1 (37)	No	
Tc-99m MAG3	10 (370)	No	
Tc-99m WBC's	5 (185)	Yes	48 hr
Tl-201	3 (111)	Yes	96 hr

\* "No" means that interruption of breast feeding need not be suggested, given the criterion of a limit of 1 mSv ED to the infant and these amounts of administered activity. "Yes" means that some interruption is required, as noted in the next column.